

# An injective $C^*$ -power series violating the Bieberbach coefficient bound

Research proof and Lean formalization

October 27, 2026

## Abstract

We give a normalized power series on  $C([0, 1], \mathbb{C})$  which converges in norm at every element and is globally injective, but whose second coefficient violates the proposed  $C^*$ -algebraic Bieberbach order bound. Its formal derivative series also converges everywhere. Injectivity follows from a continuous exponential period anchored at zero; injectivity of individual scalar fibers is not assumed.

Krishna's Conjecture 1.2 in [1] proposes the following extension of the scalar Bieberbach theorem. If  $\mathcal{A}$  is a unital  $C^*$ -algebra and  $F(u) = u + \sum_{n \geq 2} a_n u^n$  converges in norm and is injective on  $\{u : \|u\| < 1\}$ , then

$$a_n a_n^* \leq n^2 1, \quad a_n^* a_n \leq n^2 1 \quad (n \geq 2). \quad (1)$$

Conjecture 1.4 adds norm convergence of the derivative series. The next example violates both general-coefficient assertions. The separate invertible-coefficient variants are not claimed refuted.

**Theorem 1.** *Let  $\mathcal{A} = C([0, 1], \mathbb{C})$  with its supremum norm, and let  $a(t) = 8t$ . Put*

$$c_n = \frac{a^{n-1}}{n!} \quad (n \geq 1), \quad F(u) = \sum_{n \geq 1} c_n u^n. \quad (2)$$

*This series and its formal derivative series converge in the norm of  $\mathcal{A}$  for every  $u \in \mathcal{A}$ . The map  $F : \mathcal{A} \rightarrow \mathcal{A}$  is injective. Nevertheless  $c_2(1) = 4$ , so both inequalities in (1) fail at  $n = 2$ .*

*Proof.* For every  $u \in \mathcal{A}$ ,

$$\sum_{n \geq 1} \|c_n u^n\| \leq \|u\| \sum_{k \geq 0} \frac{\|au\|^k}{(k+1)!} < \infty.$$

The derivative series is dominated by an exponential series as well. Multiplication of (2) by  $a$  and reindexing the absolutely convergent exponential series gives the identity in  $\mathcal{A}$

$$aF(u) = \exp(au) - 1. \quad (3)$$

Here the exponential is the Banach-algebra exponential, whose scalar evaluation is the usual complex exponential. Evaluation of the norm-convergent series at  $t = 0$  also gives  $F(u)(0) = u(0)$ . In particular,

$$F(u)(t) = \frac{\exp(8tu(t)) - 1}{8t} \quad (t > 0).$$

Suppose  $F(u) = F(v)$ . By (3), the continuous function  $h(t) = 8t(u(t) - v(t))$  satisfies  $\exp(h(t)) = 1$  for every  $t$ , and  $h(0) = 0$ . Thus its image lies in the discrete set  $2\pi i\mathbb{Z}$ . Connectedness of  $[0, 1]$

forces  $h = 0$ . Since  $8t \neq 0$  for  $t > 0$ , we obtain  $u(t) = v(t)$  there, and the equality at 0 follows from  $F(u)(0) = F(v)(0)$ . Hence  $u = v$ .

Finally  $c_1 = 1$  and  $c_2(t) = 4t$ . Evaluation of either claimed order bound at  $t = 1$  would give  $16 \leq 4$ , a contradiction.  $\square$

**Why the scalar theorem does not apply.** The source requires injectivity of a map on the unit ball of the algebra. It does not require that every scalar fiber be univalent. A collision of two continuous inputs in this example would produce a continuous integer-valued exponential period, which must vanish because it vanishes at the endpoint. Some individual scalar fibers are noninjective. The classical scalar Bieberbach theorem is therefore not contradicted. The coefficients  $c_n$  for  $n \geq 2$  vanish at 0 and are not invertible elements of  $\mathcal{A}$ ; the source’s invertible-coefficient restrictions cannot be silently discarded.

**Formal scope and review status.** The Lean certificate has 28 declarations in four modules. It proves actual Banach-norm summability, the exponential identity, scalar evaluation, global injectivity, the literal coefficient-order violation, and closed negations of the two source assertions specialized to the genuine commutative unital  $C^*$ -algebra  $C([0, 1], \mathbb{C})$ . No convergence, injectivity or fiber-choice hypothesis remains. The endpoints are

`source_counterexample,      source_strong_counterexample`

in namespace `CStarBieberbach`. The isolated strict replay uses Lean 4.34.1 and mathlib revision

`d13f23b723b8a846827a245b89c10fc7d3f11612,`

with only `propext`, `Classical.choice` and `Quot.sound`. It checks the fixed coefficient  $8t$ ; no separately formalized theorem about all parameter values is claimed.

Independent final review passes all five gates, with bounded novelty coverage.

The source was read in the author-deposited Zenodo version (doi:10.5281/zenodo.6917925), whose text agrees with the source reading of Preprints version 1. Direct Preprints PDF retrieval remains unavailable. Acceptance additionally requires a fresh replay of the extracted proof archive. The two variants use one construction and constitute one outcome. Classical exponential-series and connectedness facts and helper declarations are not additional discoveries. No worldwide priority claim is made.

## References

- [1] K. Mahesh Krishna,  *$C^*$ -algebraic Bieberbach, Robertson, Lebedev–Milin, Zalcman, Krzyz and Corona Conjectures*, Preprints (2022), version 1, Conjectures 1.2 and 1.4, pp. 2–3. doi:10.20944/preprints202208.0140.v1.